

AD-A174 030

A REFINEMENT OF SARKOVSKII'S THEOREM(U) ARMY BALLISTIC
RESEARCH LAB ABERDEEN PROVING GROUND MD
N P BHATIA ET AL. SEP 86 BRL-TR-2762


1/1

UNCLASSIFIED

F/G 19/4


NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12



US ARMY
MATERIEL
COMMAND

AD

TECHNICAL REPORT BRL-TR-2762

AD-A174 030

A REFINEMENT OF SARKOVSKI'S THEOREM

Nam P. Bhatia
Walter O. Egerland

September 1986

DTIC
ELECTE
NOV 14 1986
S E D

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

US ARMY BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND

DTIC FILE COPY

86 11 14 068

Destroy this report when it is no longer needed.
Do not return it to the originator.

Additional copies of this report may be obtained
from the National Technical Information Service,
U. S. Department of Commerce, Springfield, Virginia
22161.

The findings in this report are not to be construed as an official
Department of the Army position, unless so designated by other
authorized documents.

The use of trade names or manufacturers' names in this report
does not constitute indorsement of any commercial product.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TABLE OF CONTENTS

I.	INTRODUCTION	Page 5
II.	ELEMENTARY LEMMAS	Page 6
III.	PRINCIPAL RESULTS	Page 7
IV.	REMARKS	Page 11
	REFERENCES	Page 13
	DISTRIBUTION LIST	Page 15

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



I. INTRODUCTION

Let $f: R \rightarrow R$ be continuous and $x_0 \in R$. The orbit of x_0 under f is defined as the set $\{x: x = f^n(x_0), n = 0, 1, \dots\}$, where, for every positive integer n , f^n is the n -th iterate of f , $f^1 = f$, and $f^0(x_0) = x_0$. We shall write $x_n := f^n(x_0)$ for a given $x_0 \in R$ and call x_1, x_2, \dots the successors of x_0 . A pre-orbit of a given $x_0 \in R$ is any (finite or infinite) sequence $x_0, x_{-1}, x_{-2}, \dots$ such that $f(x_{-n}) = x_{-(n-1)}$ for all n for which x_{-n} is defined. The points x_{-1}, x_{-2}, \dots in any such sequence are called predecessors of x_0 . A point c_0 is called critical if $f(c_0) = c_0$, i.e., a critical point of f is a fixed point of f . A periodic point x_0 of period $p > 1$ (p a positive integer) is a point for which the relations $f^p(x_0) = x_0$, $f^k(x_0) \neq x_0$, $1 \leq k < p$, hold. If x_0 is a periodic point of period p , its orbit is denoted by $(x_0, x_1, \dots, x_{p-1})$. We shall denote the k th iterate of x_0 under the function f^m by x_k^m , $k = 0, 1, \dots$. Thus $x_k^m := (f^m)^k(x_0) = x_{mk}$, and, in particular, $x_0^m = x_k^0 = x_0$ for all nonnegative integers k and m .

Definition. Let $f: R \rightarrow R$ be continuous and $x_0 \in R$. f has a loop of order n if x_0 has a pre-orbit $(x_0, x_{-1}, \dots, x_{-n})$ such that either

$$x_0 < x_{-n} < x_{-(n-1)} < \dots < x_{-2} < x_{-1}$$

or

$$x_0 > x_{-n} > x_{-(n-1)} > \dots > x_{-2} > x_{-1}.$$

f has an infinite loop if x_0 has an infinite pre-orbit $(x_0, x_{-1}, \dots, x_{-n}, \dots)$ such that either

$$x_0 < \dots < x_{-n} < x_{-(n-1)} < \dots < x_{-2} < x_{-1}$$

or

$$x_0 > \dots > x_{-n} > x_{-(n-1)} > \dots > x_{-2} > x_{-1}.$$

A loop of order $(n - 1)$ is called an n -periodic loop if $x_0 = x_{-n}$.

We adopt the following concise notation: we say property $P(k)$ holds if f has a periodic orbit of period k . Thus $P(1)$, $L(k)$, $L(=)$ mean that f has a critical point, a periodic loop of period k , an infinite loop, respectively. Similarly, $P^n(k)$, $L^n(k)$, $L^n(=)$ shall mean that f^n has a k -periodic orbit, k -periodic loop, an infinite loop, respectively.

In this notation, Sarkovskii's theorem and our refinement read as follows.

Theorem (Sarkovskii). Let $f: R \rightarrow R$ be continuous. Then

$$\begin{aligned} P(3) &\Rightarrow P(5) \Rightarrow P(7) \Rightarrow \dots \Rightarrow \\ P(2 \cdot 3) &\Rightarrow P(2 \cdot 5) \Rightarrow P(2 \cdot 7) \Rightarrow \dots \Rightarrow \\ P(2^2 \cdot 3) &\Rightarrow P(2^2 \cdot 5) \Rightarrow P(2^2 \cdot 7) \Rightarrow \dots \Rightarrow \\ \dots &\Rightarrow \\ P(2^3) &\Rightarrow P(2^2) \Rightarrow P(2) \Rightarrow P(1). \end{aligned}$$

Theorem (SR). Let $f: R \rightarrow R$ be continuous. Then

$$\begin{aligned} L(\infty) &\dots \Rightarrow L(5) \Rightarrow L(4) \Rightarrow L(3) \Leftarrow \\ P(3) &\Rightarrow P(5) \Rightarrow P(7) \Rightarrow \dots \Rightarrow \\ L^2(\infty) &\dots \Rightarrow L^2(5) \Rightarrow L^2(4) \Rightarrow L^2(3) \Leftarrow \\ P(2 \cdot 3) &\Rightarrow P(2 \cdot 5) \Rightarrow P(2 \cdot 7) \Rightarrow \dots \\ L^{2^2}(\infty) &\dots \Rightarrow L^{2^2}(5) \Rightarrow L^{2^2}(4) \Rightarrow L^{2^2}(3) \Leftarrow \\ P(2^2 \cdot 3) &\Rightarrow P(2^2 \cdot 5) \Rightarrow P(2^2 \cdot 7) \Rightarrow \dots \\ \dots &\dots \\ \dots &\Rightarrow P(2^3) \Rightarrow P(2^2) \Rightarrow P(2) \Rightarrow P(1). \end{aligned}$$

A. N. Sarkovskii obtained the fundamental result which bears now his name in his seminal paper¹ of 1964. The purpose of this paper is to prove Theorem (SR).

II. ELEMENTARY LEMMAS

It follows from the definition of a periodic loop that every three-periodic orbit is a three-periodic loop and that an $(n + 1)$ -periodic loop implies the existence of a loop of order n .

Lemma 2.1. If f has a critical point c_0 such that $c_0 < c_{-2} < c_{-1}$, f has an infinite loop satisfying

$$c_0 < \dots < c_{-n} < \dots < c_{-2} < c_{-1}.$$

The same statement holds with all inequalities reversed.

Proof. Since $f(c_{-2}) = c_{-1}$ and $c_0 < c_{-1}$, there exists $c_{-3} \in (c_0, c_{-2})$. Repeating this argument establishes the lemma.

Lemma 2.2. If f has a critical point c_0 such that $c_{-1} < c_{-3} < c_0 < c_{-2}$, c_0 has

¹ A. N. Sarkovskii, "Coexistence of cycles of a continuous map of a line into itself," Ukrain. Mat. Zh. 16 (1964), 61-71.

an infinite pre-orbit satisfying

$$c_{-1} < c_{-3} < \dots < c_0 < \dots < c_{-4} < c_{-2}.$$

In particular, f^2 has two infinite loops. The same statement holds with all inequalities reversed.

Proof. Since $f(c_{-2}) = c_{-1} < c_{-3}$ and $c_0 > c_{-3}$, there exists $c_{-4} \in (c_0, c_{-2})$, and since $f(c_{-3}) = c_{-2} > c_{-4} > c_0$, there exists $c_{-5} \in (c_{-3}, c_0)$. Repeating this argument proves the lemma.

Lemma 2.3. $P^{2^k}(n) \iff P(2^k \cdot n)$, $n, k = 1, 2, \dots$.

Proof. It suffices to show that $P^2(n) \iff P(2 \cdot n)$. If $(x_0, x_1, \dots, x_{2n-1})$ is a $2n$ -periodic orbit of f , $(x_0^2, x_1^2, \dots, x_{n-1}^2)$ is an n -periodic orbit of f^2 .

Hence $P(2 \cdot n) \implies P^2(n)$. If $(x_0^2, x_1^2, \dots, x_{n-1}^2)$ is an n -periodic orbit of f^2 , we consider the set $\{x_0, x_1, \dots, x_{2n-1}\}$, where $x_{2n} = x_n^2 = x_0^2 = x_0$. If $x_0 \neq x_k$, $k=1, 2, \dots, 2n-1$, then $C = (x_0, x_1, \dots, x_{2n-1})$ is a $2n$ -periodic orbit of f . Otherwise, there is a smallest odd k , $1 < k < 2n$, such that $x_0 = x_k$, i.e., x_0 is an odd-periodic point of f . But then, by Sarkovskii's theorem, f has periodic orbits of every even period and, therefore, in particular, a $2n$ -periodic orbit. Hence $P^2(n) \implies P(2 \cdot n)$ and the proof of the lemma is complete.

III. PRINCIPAL RESULTS

Let $C = (x_0, x_1, \dots, x_{n-1})$ be any n -periodic orbit of f . We define the subsets

$$\begin{aligned} C^+ &= \{x_i \in C: x_{i+1} > x_i\}, \quad C^- = \{x_i \in C: x_{i+1} < x_i\}, \\ D^+ &= \{x_i \in C: x_{i+2} > x_{i+1} > x_i\}, \quad D^- = \{x_i \in C: x_{i+2} < x_{i+1} < x_i\}. \end{aligned}$$

The sets C^+ and C^- are non-empty since $\min C \in C^+$ and $\max C \in C^-$. Letting further $a_0^+ = \min C^+ (= \min C)$, $b_0^+ = \max C^+$, $a_0^- = \min C^-$, and $b_0^- = \max C^- (= \max C)$, it is clear that either $a_0^+ \leq b_0^+ < a_0^- \leq b_0^-$ or $a_0^+ < a_0^- < b_0^+ < b_0^-$.

Theorem 3.1. If $a_0^+ \leq b_0^+ < a_0^- \leq b_0^-$ and $D^+ \cup D^- \neq \emptyset$, f has a critical point c_0 such that f^2 has two infinite loops $(d_0^2, d_{-1}^2, d_{-2}^2, \dots)$ and $(c_0^2, c_{-1}^2, c_{-2}^2, \dots)$ satisfying

$$d_{-1}^2 < d_{-2}^2 < \dots < d_0^2 = c_0 = c_0^2 < \dots < c_{-2}^2 < c_{-1}^2.$$

In particular, $L^2(*)$ holds.

Proof. It is sufficient to assume that $D^+ \neq \emptyset$. Then, if we let $\beta_0^+ = \max D^+$, we have

$$a_0^+ \leq \beta_0^+ < \beta_1^+ \leq b_0^+ < a_0^- \leq \beta_2^+ \leq b_0^-.$$

and conclude the existence of a critical point c_0 and a predecessor c_{-1} such that

$$a_0^+ \leq \beta_0^+ < c_{-1} < \beta_1^+ \leq b_0^+ < c_0 < a_0^- \leq b_0^-.$$

We consider now the set $E^- = \{x_i \in C^- : x_{i+1} < c_{-1}\}$. E^- is non-empty since $a_{n-1}^+ \in C^-$ and $a_n^+ = a_0^+ < c_{-1}$. Letting $r_0^- = \min E^-$, we have

$$a_0^+ \leq r_1^- < c_{-1} < b_0^+ < c_0 < a_0^- \leq r_0^- \leq b_0^-.$$

This shows that, since $c_0 > c_{-1}$ and $r_1^- < c_{-1}$, there exists a predecessor c_{-2} such that

$$a_0^+ < c_{-1} < b_0^+ < c_0 < c_{-2} < r_0^- \leq b_0^-.$$

Our construction implies that

$$(i) \text{ if } x_i \in C^+ \text{ and } x_i > c_{-1}, \text{ then } x_{i+1} \in C^-$$

$$(ii) \text{ if } x_i \in C^- \text{ and } x_i < r_0^-, \text{ then } x_{i+1} > c_{-1}.$$

Hence, there is an $x_i \in C^+$, $x_i > c_{-1}$ such that $x_{i+1} \geq r_0^-$. For otherwise we would have $b_i^+ \in (c_{-1}, r_0^-)$ for all i , contradicting the fact that C is the orbit of b_0^+ (thus $b_i^+ = a_0^+$ for some $i > 1$). We now choose $\delta_0^+ \in C^+$ such that $\delta_0^+ > c_{-1}$ and $\delta_1^+ \geq r_0^-$ to obtain

$$a_0^+ < c_{-1} < \delta_0^+ \leq b_0^+ < c_0 < c_{-2} < r_0^- \leq \delta_1^+ \leq b_0^-.$$

But this implies that we may choose a predecessor c_{-3} in the interval (c_{-1}, δ_0^+) , and hence that c_0 and its predecessors c_{-1} , c_{-2} , and c_{-3} satisfy the

inequality $c_{-1} < c_{-3} < c_0 < c_{-2}$. Appeal to Lemma 2.2 completes the proof.

Theorem 3.2. If $a_0^+ < a_0^- < b_0^+ < b_0^-$, there exist critical points d_0, c_0 of f and two infinite loops $(d_0, d_{-1}, d_{-2}, \dots)$ and $(c_0, c_{-1}, c_{-2}, \dots)$ of f satisfying

$$d_{-1} < d_{-2} < \dots < d_0 \leq c_0 < \dots < c_{-2} < c_{-1}.$$

In particular, $L(\cdot)$ holds.

Proof. We note first that there is $\alpha_0^- \in C^-$ and $\beta_0^+ \in C^+$ such that

- (i) $a_0^+ < a_0^- \leq \alpha_0^- < \beta_0^+ \leq b_0^+ < b_0^-$
- (ii) if $x_i \in C$, then $x_i \leq \alpha_0^-$ or $x_i \geq \beta_0^+$
- (iii) if $x_i \in C$ and $\alpha_0^- < x_i \leq b_0^+$, then $x_i \in C^+$
- (iv) if $x_i \in C$ and $b_0^+ < x_i \leq b_0^-$, then $x_i \in C^-$.

We now show that there are predecessors c_{-1} and c_{-2} of the critical point $c_0 \in (\alpha_0^-, \beta_0^+)$ that satisfy the inequality $c_0 < c_{-2} < c_{-1}$. The set $A^- = \{x_i \in C^- : x_i > b_0^+ \text{ and } x_{i+1} \leq \alpha_0^-\}$ is non-empty (otherwise $\beta_n^+ \geq \beta_0^+$ for all integers $n \geq 0$, a contradiction). Let $r_0^- = \min A^-$. We have $r_0^- > b_0^+$ and observe that the set $A^+ = \{x_i \in C^+ : \beta_0^+ \leq x_i \leq b_0^+ \text{ and } x_{i+1} \geq r_0^-\}$ is non-empty (since otherwise β_n^+ will satisfy $\beta_0^+ \leq \beta_n^+ < r_0^-$ for $n \geq 0$, a contradiction). We choose any $y_0^+ \in A^+$ and have

$$\alpha_0^- < c_0 < \beta_0^+ \leq y_0^+ \leq b_0^+ < r_0^- \leq b_0^-.$$

Hence

$$c_0 < c_{-2} < y_0^+ \leq b_0^+ < c_{-1} < r_0^- \leq b_0^-,$$

where the existence of c_{-1} follows from $b_1^+ > c_0$ and $r_1^- < c_0$ and that of c_{-2} from $c_0 < c_{-1}$ and $y_1^+ \geq r_0^- > c_{-1}$. The infinite loop $(c_0, c_{-1}, c_{-2}, \dots)$ satisfying $c_0 < \dots < c_{-2} < c_{-1}$ follows from Lemma 2.1. An analogous procedure locates a critical point d_0 and predecessors d_{-1}, d_{-2} such that $d_{-1} < d_{-2} < d_0 \leq c_0$, and hence an infinite loop $(d_0, d_{-1}, d_{-2}, \dots)$ satisfying $d_{-1} < d_{-2} < \dots < d_0 \leq$

c_0 . This completes the proof.

Theorem 3.3. If f has a loop of order $n \geq 3$, f has two distinct n -periodic loops. In particular, $L(n)$ holds.

Proof. Let $(x_0, x_{-1}, \dots, x_{-n})$ be a loop of order $n \geq 3$ of f such that

$$x_0 < x_{-n} < \dots < x_{-2} < x_{-1}.$$

Since there is a critical point $c_0 \in (x_{-2}, x_{-1})$, there are predecessors $c_{-1}, c_{-2}, \dots, c_{-(n-2)}$ such that

$$x_0 < x_{-n} < c_{-(n-2)} < x_{-(n-1)} < \dots < c_{-1} < x_{-2} < c_0 < x_{-1}.$$

We consider now the set

$$S = \{y_0 \in \mathbb{R} : y_n < y_0 < c_{-(n-2)} < y_1 < \dots < y_{n-3} < c_{-1} < y_{n-2} < c_0 < y_{n-1}\}.$$

The set S is non-empty since $x_{-n} \in S$ and open since f is continuous. Let (a_0, b_0) be the component of S such that $x_{-n} \in (a_0, b_0)$. Since $c_{-(n-2)} \notin S$ and $y_0 \in (a_0, b_0)$ implies $y_0 < c_{-(n-2)}$, we must have

$$- \infty < a_0 < b_0 < c_{-(n-2)}.$$

We first note that $a_0 > -\infty$. This is so because for every $y_0 \in (a_0, b_0)$ we have $y_n < y_0$ and $y_n \in f^2([c_{-1}, c_0])$, which is a compact set. Thus $y_0 \geq \min f^2([c_{-1}, c_0]) > -\infty$. This implies $a_0 \geq \min f^2([c_{-1}, c_0]) > -\infty$. Since $a_0, b_0 \notin S$ and $y_0 \in (a_0, b_0)$ implies $y_0 \in S$, we conclude by the continuity of f that

$$a_n = a_0 < c_{-(n-2)} < a_1 < \dots < a_{n-3} < c_{-1} < a_{n-2} < c_0 < a_{n-1}$$

and

$$b_n = b_0 < c_{-(n-2)} < b_1 < \dots < b_{n-3} < c_{-1} < b_{n-2} < c_0 < b_{n-1}.$$

Hence both a_0 and b_0 are n -periodic points, and since $a_0 < b_0 < b_1 < \dots < b_{n-1}$, the orbits of a_0 and b_0 are distinct. This completes the proof of the theorem.

Corollary 3.3. If f has an $(n+1)$ -periodic loop, $n \geq 3$, f has two distinct n -periodic loops.

Theorem 3.4. $L(\infty) \implies \dots \implies L(4) \implies L(3) \implies P(5) \implies P(7) \implies \dots \implies L^2(\infty)$.

Proof. If f has an infinite loop satisfying

$$x_0 < \dots < x_{-2} < x_{-1},$$

the subset $\{x_0, x_{-1}, \dots, x_{-n}\}$, $n \geq 3$, satisfies

$$x_0 < x_{-n} < \dots < x_{-2} < x_{-1},$$

and is, therefore, a loop of order n . By Theorem 3.3, $L(n)$ holds. Hence

$L(\infty) \Rightarrow L(n)$. By Corollary 3.3, $L(n) \Rightarrow L(n-1)$. The implications

$L(3) \Rightarrow P(5) \Rightarrow P(7) \Rightarrow \dots$ follow from Sarkovskii's theorem. Finally, to prove the implication $P(2n+1) \Rightarrow L^2(\infty)$ for every $n \geq 1$, we note that if $C = (x_0, x_1, \dots, x_{2n})$ is a $(2n+1)$ -periodic orbit, then $n(C^+) \neq n(C^-)$, so that the hypothesis of either Theorem 3.1 or Theorem 3.2 is satisfied. In the first case $L^2(\infty)$ holds by Theorem 3.1. In the second case, Theorem 3.2 implies that $L(\infty)$ holds, and hence that $L(3)$ holds. Now for any three-periodic orbit, the hypothesis of Theorem 3.1 holds trivially. Hence $L^2(\infty)$ holds. This completes the proof.

Corollary 3.4. $L^{2^k}(\infty) \dots \Rightarrow L^{2^k}(5) \Rightarrow L^{2^k}(3) \Rightarrow P(2^k \cdot 5) \Rightarrow P(2^k \cdot 7) \Rightarrow \dots \Rightarrow L^{2^{k+1}}(\infty)$.

Proof. This follows from Theorem 3.4 and Lemma 2.3.

Proof of Theorem (SR). Theorem (SR) follows by combining Theorem 3.4, Corollary 3.4, Lemma 2.3 and Sarkovskii's theorem.

IV. REMARKS

1. Theorem (SR) is a step in the direction of obtaining a complete refinement of Sarkovskii's theorem that takes into account the orbit types of each period n . A periodic loop is only one of the orbit types of a given period. That certain orbit types imply the existence of infinite loops is implicit in Theorem 3.2 and is strikingly illustrated by the example $f(x) = ax(1 - |x|)$. f has the four-periodic orbit $\left[\frac{1}{2}, \frac{a}{4}, \frac{1}{2}, -\frac{a}{4}\right]$, where $a \approx 4.411138875$ is given by

$$a^{-1} = \frac{1}{2} \left[\left(\frac{1}{2} + \frac{1}{2} \sqrt{\frac{59}{27}} \right)^{1/3} + \left(\frac{1}{2} - \frac{1}{2} \sqrt{\frac{59}{27}} \right)^{1/3} \right]$$

This orbit satisfies the conditions of Theorem 3.2 and hence guarantees the existence of two infinite loops as well as two periodic loops of each period $n \geq 3$ by Theorem (SR). The results of⁵ ensure for this example merely the existence of a three-periodic orbit.

2. The results in this paper offer a novel approach to detecting chaos. Most practical methods for detecting chaos rely, either implicitly or explicitly, on the existence of odd periodic orbits^{2,3,4,5}. However, lemmas 2.1 and 2.2 can be used by finding only a few predecessors of a critical point. Lemma 2.2, in particular, is independent of odd periods. A notable illustration is the example $f(x) = x^2 - s$, for which at

$$s^* = \frac{1}{3} \left[2 + \left(3\sqrt{33} + 17 \right)^{1/3} - \left(3\sqrt{33} - 17 \right)^{1/3} \right]$$

$L^2(\bullet)$ holds, with no odd period $\neq 1$ being present.

² Gyorgy Targonski, "Topics in Iteration Theory," Studia Mathematica, Skript 6, Vandenhock and Ruprecht, Göttingen and Zürich (1981).

³ Nam P. Bhatia and Walter O. Egerland, "On the Existence of Li-Yorke Points in the Theory of Chaos," Mathematics Research Report No. 84-4, Department of Mathematics, UMBC, July 1984. (To appear in Vol. 9, No. 10 of Nonlinear Analysis.)

⁴ Nam P. Bhatia and Walter O. Egerland, "Non-periodic conditions for Chaos and Snap-Back Repellers," Transactions of the Second Army Conference on Applied Mathematics and Computing, ARO Report 85-1 159-164 (Also Mathematics Research Report No. 84-5, Department of Mathematics, UMBC, September 1984).

⁵ Tien-Yien Li, Michal Misiurewicz, Giulio Pianigiani and James A. Yorke, "No Division Implies Chaos," Transactions of the American Mathematical Society, 273(1) (1982), 191-199.

REFERENCES

- [1] A. N. Sarkovskii, "Coexistence of cycles of a continuous map of a line into itself," Ukrain. Mat. Zh. 16 (1964), 61-71.
- [2] Gyorgy Targonski, "Topics in Iteration Theory," Studia Mathematica, Skript 6, Vandenhock and Ruprecht, Göttingen and Zürich (1981).
- [3] Nam P. Bhatia and Walter O. Egerland, "On the Existence of Li-Yorke Points in the Theory of Chaos," Mathematics Research Report No. 84-4, Department of Mathematics, UMBC, July 1984. (To appear in Vol. 9, No. 10 of Nonlinear Analysis.)
- [4] Nam P. Bhatia and Walter O. Egerland, "Non-periodic conditions for Chaos and Snap-Back Repellers," Transactions of the Second Army Conference on Applied Mathematics and Computing, ARO Report 85-1, 159-164 (Also Mathematics Research Report No. 84-5, Department of Mathematics, UMBC, September 1984).
- [5] Tien-Yien Li, Michal Misiurewicz, Giulio Pianigiani and James A. Yorke, "No Division Implies Chaos," Transactions of the American Mathematical Society, 273(1) (1982), 191-199.

<u>Copies</u>	<u>Organization</u>	<u>Copies</u>	<u>Organization</u>
12	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22304-6145	1	Commander U.S. Army Communications- Electronics Command ATTN: AMSEL-ED Fort Monmouth, NJ 07703
1	HQDA DAMA-ART-M Washington, D.C. 20310	1	Commander ERADCOM Technical Library ATTN: DELSD-L (Reports Section) Fort Monmouth, NJ 07703-5301
1	Commander U.S. Army Materiel Command ATTN: AMCDRA-ST 5001 Eisenhower Avenue Alexandria, VA 22333-0001	1	Commander U.S. Army Missile Command Research, Development & Engin- eering Center ATTN: AMSMI-RD Redstone Arsenal, AL 35898
1	Commander Armament R&D Center U.S. Army AMCCOM ATTN: SMCAR-TSS Dover, NJ 07801	1	Director U.S. Army Missile & Space Intelligence Center ATTN: AIAMS-YDL Redstone Arsenal, AL 35898-5500
1	Commander Armament R&D Center U.S. Army AMCCOM ATTN: SMCAR-TDC Dover, NJ 07801	1	Commander U.S. Army Tank Automotive Cmd ATTN: AMSTA-TSL Warren, MI 48397-5000
1	Director Benet Weapons Laboratory Armament R&D Center U.S. Army AMCCOM ATTN: SMCAR-LCB-TL Watervliet, NY 12189	1	Director U.S. Army TRADOC Systems Analysis Activity ATTN: ATAA-SL White Sands Missile Range, NM 88002
1	Commander U.S. Army Armament, Munitions and Chemical Command ATTN: SMCAR-ESP-L Rock Island, IL 61299	1	Commandant U.S. Army Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905
1	Commander U.S. Army Aviation Research and Development Command ATTN: AMSAV-E 4300 Goodfellow Blvd St. Louis, MO 63120	1	Commander U.S. Army Development and Employ- ment Agency ATTN: MODE-TED-SAB Fort Lewis, WA 98433
1	Director U.S. Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035	1	AFWL/SUL Kirtland AFB, NM 87117
		1	Air Force Armament Laboratory ATTN: AFATL/DLODL Eglin AFB, FL 32542-5000

CopiesOrganization

10 Central Intelligence Agency
Office of Central Reference
Dissemination Branch
Room GE-47 HQS
Washington, D.C. 20502

ABERDEEN PROVING GROUND

Dir, USAMSAA
ATTN: AMXSY-D
AMXSY-MP, H. Cohen

Cdr, USATECOM
ATTN: AMSTE-TO-F

Cdr, CRDC, AMCCOM
ATTN: SMCCR-RSP-A
SMCCR-MU
SMCCR-SPS-IL

USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. BRL Report Number _____ Date of Report _____

2. Date Report Received _____

3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) _____

4. How specifically, is the report being used? (Information source, design data, procedure, source of ideas, etc.) _____

5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided or efficiencies achieved, etc? If so, please elaborate. _____

6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) _____

CURRENT ADDRESS Name _____
 Organization _____
 Address _____
 City, State, Zip _____

7. If indicating a Change of Address or Address Correction, please provide the New or Correct Address in Block 6 above and the Old or Incorrect address below.

OLD ADDRESS Name _____
 Organization _____
 Address _____
 City, State, Zip _____

(Remove this sheet along the perforation, fold as indicated, staple or tape closed, and mail.)

----- FOLD HERE -----

Director
U.S. Army Ballistic Research Laboratory
ATTN: SLCBR-DD-T
Aberdeen Proving Ground, MD 21005-5066




NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

BUSINESS REPLY MAIL
FIRST CLASS PERMIT NO 12062 WASHINGTON, DC
POSTAGE WILL BE PAID BY DEPARTMENT OF THE ARMY

Director
U.S. Army Ballistic Research Laboratory
ATTN: SLCBR-DD-T
Aberdeen Proving Ground, MD 21005-9989



----- FOLD HERE -----

END

12-86

DTIC